

## 草莓中黄酮类物质的测定以及与抗氧化活性之间的相互关系

郑虎哲<sup>1,2</sup>, Lee Hye-ryun<sup>1</sup>, 崔春兰<sup>2</sup>, Hwang In-wook<sup>1</sup>,  
Kwon Sun-young<sup>1</sup>, Chung Shin-kyo<sup>1\*</sup>

(1 韩国国立庆北大学农学院食品工学部, 大邱 702-701, 韩国;

2 辽宁农业职业技术学院农产品质量安全检测中心, 沈阳 营口 115009)

**摘要:** 报道草莓中黄酮类物质含量的测定方法, 并比较了总抗氧化成分与抗氧化活性 (DPPH, FRAP) 之间的相互关系。黄酮类物质的水解随着 HCl 浓度和水解时间的不同发生较大的差异, 结果表明, 总多酚类物质和总黄酮类物质之间有很高的相关关系, 总多酚类物质和总黄酮类物质对 DPPH 有很高的相关关系, 而且总多酚类物质和总黄酮类物质对 FRAP 的相关关系各为  $r = 0.958$ ,  $P < 0.05$  和  $r = 0.936$ ,  $P < 0.05$ , 从结果可以推测: 草莓体现出很强的抗氧化活性, 这与草莓中的总多酚类物质成分密不可分。

**关键词:** 草莓; 水解浓度; 水解时间; 总多酚类物质; 总黄酮类物质; DPPH; FRAP

中图分类号: S 482.4 + 4

文献标识码: A

文章编号: 0253-2700 (2008) 01-125-04

## Quantification of Flavonoid Compounds from Strawberry (*Fragaria ananassa*) and Correlation Effects with Antioxidant Activities

ZHENG Hu-Zhe<sup>1,2</sup>, LEE Hye-Ryun<sup>1</sup>, CUI Chun-Lan<sup>2</sup>, HWANG In-Wook<sup>1</sup>,  
KWON Sun-Young<sup>1</sup>, CHUNG Shin-Kyo<sup>1\*</sup>

(1 Department of Food Science and Technology, Kyungpook National University, Daegu 702-701, Korea;

2 Agricultural Products Analysis Center, Liaoning Agricultural College, Yingkou 115009, China)

**Abstract:** The amounts of flavonoid (myricetin, quercetin and kaempferol), antioxidant contents (total phenolic and total flavonoid) and antioxidant activities (DPPH and FRAP) were analyzed in strawberry. Quantitative analysis of conjugated flavonoid contents were obtained after different hydrolysis concentration and reaction period. The highest yields of myricetin and quercetin were found with 1.0M HCl and a reaction period of 0.5 hr, however, for kaempferol was 1.0M HCl and a reaction period of 1hr, respectively. In the study, we also estimated the correlation coefficient of TPC, TFC, DPPH and FRAP. In addition, a positive correlation was observed ( $r = 0.980$ ,  $P < 0.05$ ) between TPC and TFC, TPC and TFC are correlated to the DPPH activity ( $r = 0.913$ ,  $P < 0.05$  and  $r = 0.899$ ,  $P < 0.05$ , respectively), and to the FRAP activity ( $r = 0.958$ ,  $P < 0.05$  and  $r = 0.936$ ,  $P < 0.05$ , respectively), respectively. These results pointed out that strawberry generally possesses a high level of antioxidant activities, which could be linked to the levels of phenolic compounds in the fruit.

**Key words:** Strawberry; Hydrolysis concentration; Reaction period; Antioxidant contents; Antioxidant activities; Correlation coefficient

Strawberry fruits, with its special sweet, sour taste and exquisite flavor, especially have previously been shown to contain antioxidant compounds such as flavonoid (myricetin, quercetin

and kaempferol) and phenolic acids, which provide protection against harmful free radicals (Vinson *et al.*, 2001, Cai *et al.*, 2004, Jose *et al.*, 2007). Flavonoids have been detected in

\* Author for correspondence; E-mail: kchung@knu.ac.kr

Received date: 2007-05-24, Accepted date: 2007-07-10

作者简介: 郑虎哲 (1974-) 男, 农学博士, 主要从事食品分析研究。

many studies with plants, measured by after acid hydrolysis, because most flavonoids founded in plants are conjugated to sugars as glycosides, hydrolysis of all glycosides to aglycones offers a practical method for the quantitative determination of flavonoids in plant (Hertog and Peter, 1992; Alan *et al.*, 1997) .

To the best of the author s knowledge, the flavonoids aglycones in strawberry have not yet been estimated according different experiment design of hydrolysis concentration and reaction period . Furthermore the correlation among antioxidant contents such as total phenolic content (TPC) , total flavonoid content (TFC) , and free radical scavenging activity of DPPH, FRAP also have not been reported . The objectives of this study were 1) to optimize the methods of extraction and hydrolysis of flavonoids conditions and 2) to determine DPPH and FRAP activity to estimate antioxidant activities and their correlations with TPC and TFC in strawberry fruit .

Materials and Methods

Strawberry ( *Fragaria ananassa* Duch .) was purchased from the local market in Dae-gu Korea .

Extraction and hydrolysis fruit for flavonoid aglycones analysis were obtained by different reaction and acid concentration . And the treated sample was dissolved with methanol and then filtered through a 0.45 μm membrane filter for HPLC analysis .

The total phenolic content (TPC) and DPPH was determined according Cai *et al.* (2004) , measurement of total flavonoid content (TFC) and FRAP was determined according to Faria *et al.* (2005) with some modifications .

Results and Discussion

We estimated the optimum extraction and hydrolysis conditions for quantifying the fruit of strawberry, because the completeness of hydrolysis largely depended on the type of glycosides . Four hydrochloric acid concentrations were tested (0.5, 1.0, 1.5 and 2.0 mol/L) , and the reaction period was varied (0.5 , 1 and 2 hrs) in the procedure described under extraction and hydrolysis . The influence of acid concentration and reaction period corresponding aglycone yield is presented in table 1 .

In our study, RP-HPLC coupled with UV-VIS Detector was employed to separate, identify and quantify favonoid aglycones in strawberry fruit ( Fig . 1 ) . In this study, the highest yields of myricetin and quercetin were found with 1.0 mol/L HCl and a reaction period of 0.5 h, however, for kaempferol was 1.0 mol/L HCl and a reaction period of 1 h, respectively . After 2 hrs with 2 mol/L HCl concentration, a loss of up to 50.3 % and 41.7% of myricetin and quircetin was observed compare to the optimum conditions, respectively ( Table 1 ) . These findings supported those of the study by Alan *et al.* (1997) who reported that

cooking will fall in quercetin and myrecitin content, boiling produced even bigger reduction in tomatoes and onions . All these results indicated that prolonged or shortened extraction and hydrolysis showed a decrease the flavonol contents in strawberry .

Table 1 Quantification data for optimization of hydrolysis conditions in strawberry fruit ( 100 mg g<sup>-1</sup> fresh weight )

HCl Conc . (mol/L)	time (h)	Myricetin	Quercetin	Kaempferol	Total
0.5	0.5	4.48	5.95	N.D <sup>a</sup>	10.43
	1	4.63	6.21	1.48	12.32
	2	4.13	6.07	1.32	11.52
1.0	0.5	5.09	6.60	1.45	13.14
	1	4.69	5.89	1.92	12.5
	2	3.31	5.34	1.12	9.77
1.5	0.5	4.89	6.12	1.55	12.56
	1	3.27	5.38	1.75	10.4
	2	2.86	3.66	1.66	8.18
2.0	0.5	3.88	5.02	1.77	10.67
	1	3.06	4.23	1.67	8.96
	2	2.56	2.75	N.D <sup>a</sup>	5.31

<sup>a</sup> N.D ., Not detected .




Fig . 1 Chromatograms of flavonoid aglycones, strawberry extracts and hydrolysis with 1.0 mol/L HCl and a reaction period of 1 h  
M: myricetin; Q: quercetin; K: kaempferol

The TPC and TFC in the strawberry fruits in methanol extraction showed the higher value of TPC ( 4575.8 ± 80.8 mg GAE/100 g F.W .) and TFC ( 430.7 ± 28.5 mg QE/100 g F.W .) compare to dichloromethane extraction of TPC ( 41.4 ± 5.3 mg GAE/100 g F.W .) and TFC ( 3.8 ± 2.7 mg QE/100 g F.W .) , respectively . Our result was agreement with those reported by Kriengsak *et al.* (2006) , who showed that TPC in methanol extraction is very higher compare to dichloromethane extraction . All these results indicate that methanol extraction is a better extraction solvent for the extraction of both TPC and TFC .

The methanol extraction exhibited a concentration-dependent




Fig. 2 Free radical scavenging activity (DPPH) of strawberry means  $\pm$  SD,  $n = 3$ . Bars represent standard error




Fig. 3 Ferric reducing antioxidant power (FRAP) of strawberry means  $\pm$  SD,  $n = 3$ . Bars represent standard error

DPPH radical scavenging activity, which was higher than that of the positive control, trolox, the scavenging effect increased with the increasing concentrations from 0.8 - 8.0 mg/ml, it was found to be over 90.8% in 8.0 mg/ml concentration, which is significantly higher than that of gallic acid (82.9%) and trolox (30.4%) at the same concentration of 150  $\mu$ mol (Figure 2). These results were agreement with Jose *et al.* (2007) who indicated that native cultivated strawberries (*F. vesca*) were 2.5 times more active than in the TEAC assay (Trolox equivalent antioxidant capacity).

Figure 3 clearly showed that the FRAP of methanol extract of strawberry had reducing power of  $191.5 \pm 7.2$  TE  $\mu$ mol/mg extract. This was slight higher than that of ascorbic acid ( $184.5 \pm 5.6$  TE  $\mu$ mol/mg), but lower than that of gallic acid ( $447.7 \pm 5.9$  TE  $\mu$ mol/mg) within a concentration of 150  $\mu$ m. According to its high value, it could be considered that compound was good electron donors and could terminate oxidation chain reactions by reducing the oxidized intermediates into the stable form.

Recently studies have suggested that the antioxidant contents and antioxidant free radical scavenging activities have the positive effect (Jose *et al.*, 2007, Kriengsak *et al.*, 2006). In our study with the strawberry fruit, a positive correlation was observed ( $r = 0.980$ ,  $P < 0.05$ ) between TPC and TFC in methanol extraction, however, there was no significant correlation in dichloromethane extraction (data not shown). Furthermore, TPC and TFC are correlated to the DPPH activity ( $r = 0.913$ ,  $P < 0.05$  and  $r = 0.899$ ,  $P < 0.05$ , respectively) and to the FRAP activity ( $r = 0.958$ ,  $P < 0.05$  and  $r = 0.936$ ,  $P < 0.05$ , respectively) was observed. In addition, a positive correlation was also observed ( $r = 0.947$ ,  $P < 0.05$ ) between DPPH and

FRAP. Our results indicated that TPC showed the highest correlation with TFC. It could be explain that phenolic and flavonoid compounds, which are known as hydrophilic antioxidants, are secondary metabolites that are most abundant in fruits, which positively correlated with the oxygen radical absorbance capacity (ORAC). In addition, most techniques used for determining antioxidant activity, showed high correlation with TPC and TFC in different free radical scavenging activities. All these results were agreement with Vinson *et al.* (2001) who reported that strawberry generally possesses a high level of antioxidant activity, which could be linked to the levels of phenolic compounds in the fruit. Furthermore, our results were supported by Gil *et al.* (2002), who found high correlation between antioxidant activities and total phenolic content in nectarines, peaches and plums, and they demonstrated that there was no correlation between ascorbic acid and antioxidant activity as determined by DPPH or FRAP assays.

Strawberry fruits contain a group of natural antioxidants that could have not only a high antioxidant activity but also a good combination or mixture of antioxidants, the supplementation of these natural antioxidants through a balanced diet containing enough fruits could be much more effective and also economical than the supplementation of synthetic preservatives.

## References:

- Alan Crozier, Michael EJL, Morag SM *et al.*, 1997. Quantitative analysis of the flavonoid content of commercial tomatoes, onions, lettuce, and celery [J]. *J Agric Food Chem*, **45**: 590—595
- Cai YZ, Luo Q, Sun M *et al.*, 2004. Antioxidant activity and phenolic compounds of 112 Chinese medicinal plants associated with anticancer [J]. *Life Sci*, **74**: 2157—2184
- Faria A, Oliveira J, Neves P *et al.*, 2005. Antioxidant properties of pre-

- pared blueberry (*Vaccinium myrtillus*) extracts [J]. *J Agric Food Chem*, **53**: 6896—6902
- Gil MI, Tomas-Barberan FA, Hess-Pierce B *et al.*, 2002. Antioxidant capacities, phenolic compounds, carotenoids, and vitamin C contents of nectarine, peach, and plum cultivars from *California* [J]. *J Agric Food Chem*, **50**: 4976—4982
- Hertog Michael GL, Peter CHH, 1992. Optimization of a quantitative HPLC determination of potentially anticarcinogenic flavonoids in vegetables and fruits [J]. *J Agric Food Chem*, **40**: 1591—1598
- Jose C, Cristina T b, Jaime AR b *et al.*, 2007. Free radical scavenging activity and phenolic content in achenes and thalamus from *Fragaria chiloensis* ssp. *chiloensis*, *F. vesca* and *F. xananassa* cv. Chandler [J]. *Food Chem*, **102**: 36—44
- Kriengsak T, Unaroj B, Kevin C *et al.*, 2006. Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts [J]. *J Food Compos Anal*, **19**: 669—675
- Vinson JA, Su X, Zubik L *et al.*, 2001. Phenol antioxidant quantity and quality in foods: Fruits [J]. *J Agric Food Chem*, **49**: 5315—5321

\* \* \* \* \*

〔上接 114 页〕

C. 文献排列顺序：中文、英文。中文以姓氏汉语拼音字母顺序为序，英文以姓氏字母为序（包括用汉语拼音所列作者文献）。

2.6 对植物系统学与生物地理学稿件的要求：（1）报道个别省级新分布或新记录种的稿件和描述单个新变种的稿件不接受发表，报道二个及以上中国新记录种的稿件可酌情考虑以用简报的形式发表，但篇幅不超过一个页码；（2）关于单个种的核型分析稿件不再接受发表，单种属的核型研究论文，内容充实者和确有新意者可接受投稿。

2.7 对植物生态学与资源管理稿件的要求：（1）植物生态学稿件：要求在系统综述国内外前期研究工作的基础上，明确提出科学假设（说），制定与此相对应的研究方案，通过野外调查（监测）、实验室控制实验等方法采集数据，在统计学分析结果的支持下，验证作者提出的科学假设，以解决关键科学问题，为植物生态学的理论发展做出新贡献。没有具体科学目标的、描述性的植物群落学调查资料原则上不再接受发表。（2）植物资源管理稿件：研究对象主要控制在物种、生态系统和景观水平。稿件分为两类：第一类是应用生态学的基本原理，研究或构建具有良好经济效益和生态效益的植物资源管理体系，并分析其生态系统的功能和维持机制；第二类是新开发的植物资源报道或系统整理（包括物种 亚种 变种 品种、群落等），应该明确说明资源的现状（分布或储量）、用途、开发利用前景等。第二类稿件一般作为研究简报发表，篇幅不超过 4 个印刷页。

2.8 对于植物生理与分子生物学稿件的要求：要求论文的内容对认识植物具有重要的或新的生物学事件、过程的意义、功能有贡献，不再接受仅仅描述配方改进、实验过程调整和测量数据堆砌的论文。

2.9 对植物化学稿件的几点要求：（1）原则上论文请用英文撰写；（2）论文中要有新化合物；（3）或已知化合物新的生物活性；（4）若无新化合物，但前人对其化学成分缺乏研究，可以用简报的形式发表，但篇幅不超过一个页码，主要说明其中各种成分即可，不必罗列大量分析数据和化学结构。

### 3 特别说明

3.1 鼓励通过本刊网上投稿系统（<http://journal.kib.ac.cn>）投稿，也可通过 Email 发电子文件进行投稿，稿件要求采用 Word 文档格式，具体投稿要求和格式可从本刊网站下载。

3.2 综述性论文主要以主编或学科副主编约稿的形式获得，并经审稿同意后发表。

3.3 为了加快稿件周期，本刊在审稿程序中设有绿色通道，学科副主编将在收到稿件的第一时间根据自己对稿件内容涉及的学科领域、稿件质量的判断，提出稿件进入 1) 正常审稿程序、2) 退稿或 3) 进入绿色通道（即直接向主编推荐接受，优先刊登）的建议。稿件的录用与否一般会在收到来稿 2 个月内及时给予通知。印刷稿不退，请自留底稿及原图。

3.4 为使作者的稿件顺利通过审稿和及时发表，请作者在撰写稿件时严格按我刊格式准备，写作时做到思路清晰、逻辑严密、句子通顺、衔接过渡、前后呼应、凸显创新、详简适度。对格式不符、英文摘要不地道、输入文字错误较多、句子不通或思路不清晰的稿件，编辑部将直接退稿，待作者完善了相应部分后再进入审稿程序。

### 4 联系方式

地址：昆明市黑龙潭昆明植物研究所内，《云南植物研究》编辑部 邮政编码：650204

电话：(0871) 5223032；传真：(0871) 5223032；E-mail: [bianji@mail.kib.ac.cn](mailto:bianji@mail.kib.ac.cn), <http://journal.kib.ac.cn>